



Pfizer Australia Pty Ltd  
 ABN 50 008 422 348  
 38-42 Wharf Road  
 West Ryde NSW 2114

# Boys in utero 24-42 weeks & post natal 0-3 years

Surname \_\_\_\_\_

Identification No. \_\_\_\_\_


Given Names \_\_\_\_\_

Date of Birth \_\_\_\_\_

## Head Circumference


**Measuring Technique:** The tape should be placed over the eyebrows, above the ears and over the most prominent part of the occiput taking a direct route. A paper tape is preferable to plastic, which stretches unacceptably under tension. The maximum measurement should be recorded to the nearest 0.1 cm.

### In utero 28-40 weeks, 0-12 months



Data Source: Head circumference 28-40 weeks gestation from Kitchen, W.H. *Aust. Paediatr. J.* (1983) 19:157-161).  
 Head circumference 0-3 years from [www.cdc.gov/nchs/about/major/nhanes/growthcharts/datafiles.htm](http://www.cdc.gov/nchs/about/major/nhanes/growthcharts/datafiles.htm)

### 1-3 years



Data Source: [www.cdc.gov/nchs/about/major/nhanes/growthcharts/datafiles.htm](http://www.cdc.gov/nchs/about/major/nhanes/growthcharts/datafiles.htm)

# Length Percentile for Boys 0-3 years

Mother's Height


Father's Height

**Supine Length** (recommended up to the age of 3 so that there is overlap with standing height at 2 to 3) is taken on a flat surface, with the child lying on his back. One observer holds the child's head in contact with a board at the top of the table and another straightens the legs and turns the feet upward to be at right angles to the legs and brings a sliding board in contact with the child's heels.

$$C = M[1 + L.S.Z]^{1/L}$$

Where C is the centile required, LMS are those parameters published by CDC and Z is the standard deviation equivalent to the centile required.

1st Centile calculated by Associate Professor Peter Davies, Children's Nutrition Research Centre, Brisbane.







# Boys in utero 24 - 42 weeks & post natal 0 - 3 years

## Intrauterine Growth Curves (Composite Male/Female)

**Measuring Technique:** As for ages 0-36 months (see previous pages).

**Additional Notes:** Gestational ages are recorded in completed weeks from the first day of the mother's last menstrual period. Foetal growth is influenced by many factors including age, body weight, height, parity, ethnic origin of the mother and sex of the foetus. Corrections for some of these factors are found in the quoted reference.

Data Source: Kitchen, W.H. et al. 1983, 'Revised intrauterine growth curves for an Australian hospital population', *Aust. Paediatr. J.* 19:157-161.



Endorsed By  
**A**ustralasian **P**aediatric **E**ndocrine **G**roup

The opinions, views and recommendations expressed in this publication do not necessarily reflect those of the sponsor or publisher. Pfizer Australia accepts no responsibility for treatment decisions based upon these charts.

To reorder, please call Pfizer Customer Service 1800 629 921.